

Formation of molecules near a Feshbach resonance in a 1D optical lattice

G. Orso,¹ L.P. Pitaevskii,^{1,2} S. Stringari,^{1,3} and M. Wouters^{1,4}

¹*Dipartimento di Fisica, Università di Trento and BEC-INFM, 1-38050 Povo, Italy*

²*Kapitza Institute for Physical Problems, 117334 Moscow, Russia*

³*École Normale Supérieure and Collège de France,*

Laboratoire Kastler Brossel, 24 Rue Lhomond 75231 Paris, France

⁴*TFVS, Universiteit Antwerpen, Universiteitsplein 1, 2610 Antwerpen, Belgium*

We calculate the binding energy of two atoms interacting near a Feshbach resonance in the presence of a 1D periodic potential. The critical value of the scattering length needed to produce a molecule as well as the value of the molecular binding energy in the unitarity limit of infinite scattering length are calculated as a function of the intensity of the laser field generating the periodic potential. The Bloch bandwidth and the effective mass of molecules are shown to depend strongly on the value of the scattering length due to the correlated motion of the two atoms.

PACS numbers:

The availability of tunable periodic potentials of optical nature (see, for example, Ref.[1]) and the quantum manipulation of the interatomic forces via Feshbach resonances [2] are opening new perspectives in the field of ultracold atomic gases. In the absence of the external potential molecules can be formed near resonance only for positive values of the s-wave scattering length a , the corresponding binding energy being given by the formula $E_b = -\hbar^2/ma^2$ where m is the mass of each atom. These molecules have already been observed experimentally by several groups [3, 4, 5, 6]. An interesting and very rich problem is the formation of molecules in an optical lattice generated by laser fields. Recently, Fedichev et al. [7] investigated the case of a 3D tight lattice and showed that the bound state can be formed also for negative scattering lengths down to a critical value. Their approach is valid provided that the binding energy is small compared to the width of the lowest Bloch band so that the molecular wavefunction extends over many lattice sites.

In this Letter we discuss the formation of molecules in a 1D optical lattice for *arbitrary* values of the scattering length and the laser intensity. If the barriers separating two consecutive wells are very high (high laser intensity) the system behaves like a series of 2D discs which lie at the bottom of the wells and feel a harmonic potential in the direction of the laser field. In this asymptotic regime the two-body problem is significantly simplified by the possibility of separating the center of mass and the relative coordinates of the two interacting atoms [8]. When this condition is violated many interesting questions can be addressed: what is the critical value of the scattering length required to ensure the formation of a molecule? What is the value of the binding energy of the molecule in the so called unitarity limit where the scattering length approaches an infinite value? How do the tunneling properties (bandwidth and effective mass) of molecules depend on the value of the scattering length? The purpose of this Letter is to provide an explicit answer to these questions.

Let us start by writing the 3D Schrodinger equation for two atoms interacting in the presence of a 1D periodic potential $V_{opt}(z)$:

$$\left(-\frac{\hbar^2}{m}\nabla^2 - \frac{\hbar^2}{4m}\frac{\partial^2}{\partial Z^2} + V(Z, z) + g\delta(\mathbf{r})\frac{\partial}{\partial r}\right)\Psi = E\Psi, \quad (1)$$

where $V(Z, z) = V_{opt}(z_1) + V_{opt}(z_2)$ and we have introduced the center of mass and relative coordinates $Z = (z_1 + z_2)/2$ and $\mathbf{r} = \mathbf{r}_1 - \mathbf{r}_2$.

In Eq.(1) the interaction is modeled via a *s*-wave pseudopotential with coupling constant $g = 4\pi\hbar^2 a/m$, where a is the scattering length in vacuum. The solution of Eq.(1) can be written, for $g \neq 0$, as

$$\Psi(\mathbf{r}, Z) = \int dZ' G_E(\mathbf{r}, Z; \mathbf{0}, Z') g \frac{\partial}{\partial r'} (r' \Psi(\mathbf{r}', Z'))_{\mathbf{r}'=0}, \quad (2)$$

where G_E is the free ($g = 0$) Green's function associated to Eq.(1). At short distance, the Green's function is determined only by the kinetic energy. Neglecting the external potential and setting $E = 0$ in Eq.(1), we find $G_{E=0}(\mathbf{r}, Z; \mathbf{0}, Z') = -(m/8\pi^2\hbar^2)[(Z - Z')^2 + (r/2)^2]^{-1}$. Therefore the Green's function has the following expansion

$$G_E(\mathbf{r}, Z; \mathbf{0}, Z') = -\frac{m}{4\pi\hbar^2 r} \delta(Z - Z') + K_E(Z, Z') + O(r), \quad (3)$$

where the regular kernel $K_E(Z, Z')$ depends on the external potential and we used the representation of the delta function $\delta(x) = \lim_{r \rightarrow 0} r/\pi(r^2 + x^2)$. When inserted into Eq.(2), Eq.(3) implies, for $r \rightarrow 0$, the boundary condition $\Psi(\mathbf{r}, Z) \sim (1/r - 1/a)f(Z)$, where $f(Z)$ is a function of the center of mass position satisfying the integral equation

$$\frac{1}{g}f(Z) = \int dZ' K_E(Z, Z') f(Z'). \quad (4)$$

If the center of mass and the relative motion decouple the function $f(Z)$ must be an eigenstate of the center of

mass Hamiltonian independent of the scattering length. By inserting it into Eq.(4), the dependence on Z factors out and we are left with an algebraic equation yielding the energy as a function of the scattering length. For instance, in the absence of the optical lattice the kernel is given by

$$K_E^0(Z, Z') = \frac{m}{4\pi\hbar^2} \int \frac{dP}{2\pi} e^{iP(Z-Z')} \sqrt{|E| + \hbar^2 P^2/4m}. \quad (5)$$

The lowest energy solution of Eq.(4) corresponds to $f(Z)$ constant and yields $E = -\hbar^2/ma^2$.

For periodic potentials, the kernel has the property $K_E(Z, Z') = K_E(Z + jd, Z' + jd)$ where j is any integer and d is the lattice spacing. This means that the general solution of Eq.(4) has the Bloch form $f(Z) = e^{iQZ} f_Q(Z)$, where $f_Q(Z)$ is a periodic function, Q being the quasi-momentum of the molecule. In the following we shall restrict to the lowest bound state given by Eq.(4). The energy as a function of the scattering length is then fixed by the condition that the lowest eigenvalue of the kernel $K_E(Z, Z')$ in Eq.(4) is equal to $1/g$.

Due to the singular term in Eq.(3), the numerical evaluation of the kernel $K_E(Z, Z')$ from the Green's function G_E is not trivial. The latter can be written in the general form

$$G_E(\mathbf{r}, Z; \mathbf{0}, Z') = \sum_{n_1, n_2} \int \frac{d^2 \mathbf{k}_\perp}{(2\pi)^2} \int_{-q_B}^{q_B} \frac{dq_1}{2\pi} \frac{dq_2}{2\pi} e^{i\mathbf{k}_\perp \cdot \mathbf{r}} \frac{\phi_{n_1, q_1}(Z) \phi_{n_2, q_2}(Z) \phi_{n_1, q_1}^*(Z') \phi_{n_2, q_2}^*(Z')}{E - \epsilon_{n_1}(q_1) - \epsilon_{n_2}(q_2) - \hbar^2 k_\perp^2/m}, \quad (6)$$

where $q_B = \pi/d$ is the Bragg wave-vector and $\phi_{n, q}(z_1)$ are the eigenstates of $H_z = -(\hbar^2/2m)d^2/dz_1^2 + V_{opt}(z_1)$ with energies $\epsilon_n(q)$. In the following the periodic potential will be taken of the form

$$V_{opt}(z_1) = sE_R \sin^2\left(\frac{\pi z_1}{d}\right), \quad (7)$$

as usually provided by two counterpropagating laser beams (optical lattices), where s is the laser intensity, the spacing d being related to the wavelength of the laser fields and $E_R = \hbar^2 \pi^2 / 2md^2$ is the recoil energy. Since the singular term in the rhs of Eq.(3) does not depend on the external potential, we add and subtract from G_E , the Green function (6) evaluated for $s = 0$. This permits us to write the kernel as $K_E(Z, Z') = \lim_{r \rightarrow 0} [G_E(\mathbf{r}, Z; \mathbf{0}, Z') - G_E^0(\mathbf{r}, Z; \mathbf{0}, Z')] + K_E^0(Z, Z')$, with K_E^0 defined in Eq.(5). After these manipulations, the integration over $\mathbf{k}_\perp$ in Eq.(6) converges and can be performed analytically. The remaining summation over band indices and the integration over quasi-momentum are done numerically. We define the binding energy as $E_b(Q) = E(Q) - E_{ref}(Q)$, where $E_{ref}(Q) = 2\epsilon_1(Q/2)$ is the lowest energy state for two non interacting particles with total quasi-momentum Q




FIG. 1: Binding energy versus inverse scattering length for different values of the laser intensity: from top to bottom $s = 20, 10, 5, 0$ (solid line). Also shown are the asymptotic behaviour for large d/a (dotted line) and the binding energy in harmonic approximation (dashed line) for $s = 20$.




FIG. 2: Critical value of the inverse scattering length as a function of the laser intensity.

and ϵ_1 is the dispersion of the lowest Bloch band. With this notation, E_b is always negative for a bound state.

The binding energy for quasimomentum $Q = 0$ as a function of the inverse scattering length is shown in Fig.1 for different values of the laser intensity.

When the binding energy is large with respect to the depth of the lattice ($\hbar^2/ma^2 \gg sE_R$), the periodic potential (7) can be treated as a perturbation and the wavefunction takes the form $\Psi(\mathbf{r}, Z) = \varphi(Z)\phi_b(r)$, with $\phi_b(r) = (1/\sqrt{2\pi a})e^{-r/a}/r$. Within first order perturbation theory, the effective potential acting on the center of mass of the molecule is given by $U_{eff}(Z) = \int \phi_b^2(r) V(Z, z) d^3 \mathbf{r}$, yielding

$$U_{eff}(Z) = sE_R - sE_R \frac{2}{aq_B} \arctan\left(\frac{aq_B}{2}\right) \cos(2q_B Z). \quad (8)$$

We see from Eq.(8) that the depth of the effective potential depends on the scattering length. In the limit $a \ll d$,




FIG. 3: Binding energy at unitarity ($d/a = 0$) as a function of the laser intensity (solid line). Also shown are the harmonic approximation (dashed line) and the inclusion of anharmonic corrections (solid circles).

the external potential is constant on the scale of the size of the molecule and one finds $U_{eff}(Z) = 2V_{opt}(Z)$. When a increases, the size of the molecule becomes comparable to the lattice spacing and the effective potential (8) is reduced. For $\hbar^2/ma^2 \gg sE_R$, the binding energy is given by

$$E_b(Q) = -\hbar^2/ma^2 + E'(Q) - E_{ref}(Q), \quad (9)$$

where E' can be found by solving the Schrodinger equation

$$\left[-\frac{\hbar^2}{4m} \nabla_z^2 + U_{eff}(Z) \right] \varphi(Z) = E' \varphi(Z). \quad (10)$$

The asymptotic behaviour based in Eqs (8)-(10) is plotted in Fig.1 with dotted line for different value of s .

When the laser intensity becomes large ($s \gg 1$) and the binding energy is large compared to the width of the lowest Bloch band, the system enters the quasi-2D regime. In this limit, the two interacting atoms are localized at the bottom of the same optical well where, to a first approximation, the potential (7) is harmonic with frequency $\omega_0 = 2\sqrt{s}E_R/\hbar$. In this case, the center of mass and the relative motion decouple and the scattering problem can be solved analytically [8, 9]. The corresponding result for the binding energy as a function of the inverse scattering length is shown in Fig.1 for $s = 20$ (dashed line). We see that the harmonic approximation slightly overestimates the correct binding energy.

An important consequence of the presence of the optical lattice is the formation of a bound state also for negative scattering length. However, unlike in the asymptotic 2D limit, the bound state disappears for values of the inverse scattering length below a critical value d/a_{cr} . The dependence of d/a_{cr} as a function of the laser intensity is shown in Fig.2. We see that $d/a_{cr} = 0$ for $s = 0$ and $d/a_{cr} \rightarrow -\infty$ as $s \rightarrow \infty$.

The behaviour of the binding energy as a function of the scattering length sufficiently close to the critical point is of the form $1/a - 1/a_{cr} = \sqrt{|E_b|m^*/\hbar C}$, where m^* is the atomic effective mass evaluated at the bottom of the lowest Bloch band and $C = C(s)$ is a dimensionless function of the laser intensity. For $s = 0$ one has $C = 1$ while in tight binding limit one finds $C = d/\sqrt{2\pi}\sigma$, where σ is the width of the variational gaussian ansatz $w(z) = \exp(-z^2/2\sigma^2)/\pi^{1/4}\sigma^{1/2}$ for the Bloch function $\phi_{1,q}(z) \sim \sum_{\ell} e^{iq\ell d} w(z - \ell d)$. This is just the renormalization factor of the coupling constant g obtained by integration over the gaussian profile [10]. The above results show that the optical lattice gives rise to an effective shift of the resonance, in analogy with the case of a 3D tight optical lattice investigated in Ref.[7].

It is interesting to discuss explicitly the behaviour of the binding energy at resonance ($1/a = 0$) as a function of the laser intensity. The numerical result is shown in Fig.3 (solid line). For large s the harmonic approximation gives $E_b^{ho}(1/a = 0) = -0.244\hbar\omega_0 = -0.488\sqrt{s}E_R$ (dashed line). The remaining shift ΔE in the binding is due to the non harmonic terms. We have found that the inclusion of such corrections via first order perturbation theory results in an s -independent shift $\Delta E = 0.198E_R$ which well reproduces the full numerical result for $s \gtrsim 10$ (see dots in Fig.3).

So far we have discussed the results for the binding energy corresponding to vanishing values of the quasi-momentum. Differently from the case of harmonic trapping, where the binding energy does not depend on the center of mass motion, in the presence of an optical lattice the binding energy of the molecule depends on its quasi-momentum. This is plotted in Fig.4 for $s = 2.5$ and different value of the scattering length. We see that the modulus of the binding energy increases when Q increases. Moreover, when the scattering length is negative and d/a crosses the critical value d/a_{cr} , the bound state with zero quasi-momentum breaks first. The bandwidth, defined as $w = E(q_B) - E(0)$, is plotted in Fig.5 (Left Panel) as a function of d/a for $s = 2.5$. For a given value of the laser intensity, the curve stops at $d/a = d/a_{cr}$ where $w = 2\epsilon_1(q_B/2) + E_b(q_B) < 2\epsilon_1(q_B/2)$ because the binding energy at $Q = q_B$ is finite at the critical point (see Fig.4). The bandwidth for molecules is always smaller than the one for single atoms. For instance for $s = 2.5$ the width of the lowest band for atoms is $0.52E_R$. This effect results from the correlated motion of the two constituent atoms. This is best seen for $\hbar^2/ma^2 \gg sE_R$, where the depth of the effective potential (8) felt by the molecule increases as d/a increases. In the limit $d/a \gg 1$, $U_{eff}(Z) = 4sE_{Rm} \cos^2(\pi Z/d)$, where $E_{Rm} = E_R/2$ is the recoil energy for a particle with mass $2m$, yielding an effective laser intensity $s_m = 4s$. In particular, in tight binding approximation the atomic bandwidth is proportional to the WKB tunneling exponent e^{-S_0} , where $S_0 = \pi^2 s^{1/2}/4$, while the bandwidth for




FIG. 4: Binding energy dispersion as a function of quasi-momentum Q for fixed $s = 2.5$ and for different values of the inverse scattering length: from top to bottom $d/a = 2, 0, -1.17(= d/a_{cr}), -1.66$. The solid squares give the dispersion at $d/a = 2$ evaluated from Eqs (8)-(10).




FIG. 5: Bandwidth (Left Panel) and effective mass ratio $2m/M^*$ (Right Panel) of the molecule as a function of the inverse scattering length for $s = 2.5$ (solid line). The bandwidth at the critical point is always smaller than $2\epsilon_1(q_B/2)$ (dot-dashed line) [See text]. Conversely, at the critical point the ratio $2m/M^*$ coincides with the effective mass ratio m/m^* for single atoms (dot-dashed line). The asymptotic behaviour for large positive d/a is also shown (dotted line).

molecules is much smaller, being proportional to e^{-2S_0} .

In Fig.5 (Right Panel) we show the results for the effective mass of the molecule, defined through $1/M^* = (\partial^2 E(Q)/\partial Q^2)|_{Q=0}$, as a function of d/a for laser intensity $s = 2.5$. We see that, in complete analogy with the bandwidth, the ratio $2m/M^*$ decreases as d/a increases. At the critical point where the molecule breaks, one finds $M^* = 2m^*$, where m^* is the effective mass of a single atom.

In the last part of the Letter we discuss the behaviour of the size of the molecule as a function of the scattering length and the laser intensity. The typical size of the

molecule in the radial direction is fixed by the binding energy according to $\lambda_\perp \approx \hbar/\sqrt{|E_b|m}$, where E_b is fixed by the scattering length [see Fig.(1)]. Due to the optical confinement, the molecule is more squeezed in the axial direction, so generally $\lambda_z \leq \lambda_\perp$. For large d/a the molecule is not distorted by the optical lattice and therefore $\lambda_\perp \approx \lambda_z \approx a$. When d/a decreases, the wavefunction becomes more elongated in the radial direction. As long as the binding energy is large compared to the width of the lowest Bloch band, the two atoms stay in the same optical well and therefore $\lambda_z \lesssim d$. When this condition is violated, the two atoms can hop to different lattice sites and the wavefunction becomes spread also in the z direction. Finally, close to the critical point, one has $\lambda_\perp, \lambda_z \gg d$ with the anisotropy ratio fixed by the simple relation $\lambda_\perp/\lambda_z = \sqrt{m^*/m}$.

In conclusion, we have presented an exact numerical method to calculate the binding energy and the tunneling properties of a molecule formed by two atoms interacting via a short range potential, in the presence of a 1D optical lattice, for arbitrary values of laser intensity, scattering length and quasi-momentum of the molecule. The theoretical predictions could be checked in experiments on dilute ultracold gases. They could also have important consequences on the many-body properties of these systems. In particular the formation of molecules induced by the periodic optical field is expected to modify the behaviour of the BEC-BCS crossover in a degenerate Fermi gas.

We acknowledge interesting discussions with Z. Idziaszek and C. Salomon. This work was supported by the Ministero dell'Istruzione, dell'Universita' e della Ricerca (M.I.U.R.), by the Special Research Fund of the University of Antwerp, BOF NOI UA 2004, and by the FWO-V project No.G.0435.03. M.W. is financially supported by the 'FWO - Vlaanderen'.

-
- [1] Arimondo, E., W. D. Phillips, and F. Strumia, 1992, Eds., Laser Manipulation of Atoms and Ions, Proceedings of the International School of Physics Enrico Fermi, Course CXVIII (North-Holland, Amsterdam).
 - [2] S. Inouye et al., Nature **392**, 151 (1998).
 - [3] M. Greiner, C. Regal, and D.S. Jin, Nature **426**, 537 (2003).
 - [4] S. Jochim et al., Phys. Rev. Lett. **91**, 240402 (2003).
 - [5] M. W. Zwierlein, et al., Phys. Rev. Lett. **91**, 250401 (2003).
 - [6] J. Cubizolles et al., Phys. Rev. Lett. **91**, 240401 (2003).
 - [7] P. O. Fedichev, M. J. Bijlsma, and P. Zoller, Phys. Rev. Lett. **92**, 080401 (2004).
 - [8] D.S. Petrov, M.Holzmann and G.V. Shlyapnikov, Phys. Rev. Lett. **84**, 2551 (2000); D.S. Petrov and G.V. Shlyapnikov, Phys. Rev. **A64**, 012706 (2000).
 - [9] Z. Idziaszek and T. Calarco, quant-ph/0410163. See also E.L. Bolda et al., Phys. Rev. A **68**, 032702 (2003).

- [10] P. Pedri et al., Phys. Rev. Lett. **87**, 220401 (2001).